

Risk-Assessment Worksheet

Issue: Off-label use of one minute contact times for low-level surface disinfectants used on noncritical environmental surfaces and noncritical patient care equipment.

Assessment Date: April 13, 2016

Scoring: Low = 1 Moderate = 3 High = 5

Team Members: Bill Rutala, David Weber, Becky Brooks, Lisa Teal, Emily Sickbert-Bennett, Maria Gergen, Kate Schultz, Judie Bringhurst, Elizabeth Walters

Meeting Actions: Team members evaluated the evidence and determined that off-label 1-minute contact times was sufficient to disinfect noncritical environmental surfaces and noncritical patient care equipment in a healthcare environment.

Suggested Questions	Benefit	Risk
What is the impact on patient care delivery?	There are no data demonstrating a benefit of a 10-minute contact time for surface disinfection. More than a dozen articles* demonstrate the ability of EPA-registered disinfectants to inactivate healthcare-associated pathogens (e.g., MRSA, VRE, CONS) with a contact time of ≥ 1 minutes. Additionally, data demonstrate that our surface disinfectant (QUAT) continues to have significant antimicrobial activity that extends beyond the wet time on a surface. That is, our surface disinfectant has sustained antimicrobial effectiveness (i.e., >6 hours) against HA pathogens when left on the surface. Score - 5	There is believed to be no risk to utilizing a 1-minute contact time. Score – 1
How does the issue affect the staff?	In order to achieve a contact time of 10 minutes, as recommended by the labeling on many disinfectants, requires the reapplication of the disinfectant up to	Requiring staff to follow label directions for actions with no proven benefit to employee or patient safety may serve to reduce efforts proven to improve patient



	5-6 times for a given surface, as the typical drying time of a water-based disinfectant is 1.5-2 minutes. Using a 1 minute contact time will result in more staff time available for patient care. Score = 5	outcomes. Score - 1
What is the impact on HAIs	Multiple scientific studies have demonstrated the efficacy of hospital disinfectants against pathogens causing HA infections (HAIs) with a contact time of at least 1 minute. Score - 5	There are no data that demonstrate improved infection prevention with a 10-minute contact time vs. a 1-minute contact time. Score - 1
How does the issue affect any visitors, volunteers, and so forth?	NA	NA
What is the impact on public safety?	NA	NA
What is the financial impact of the issue on the organization?	The use of a 1-minute contact time for non-critical surfaces results in more staff time spent caring for patients and less exposure to the deleterious effects of the disinfectants. Score - 3	There are no EPA enforcement action on healthcare facilities for "off-label" use of surface disinfectants. There have been citations reported from regulatory agencies such as CMS. Score - 3
What is the impact on the physical structure, including buildings, departments, units, or other areas?	NA	NA
Does the issue affect the exterior environment, including access, exit from buildings, grounds, rest areas, and so forth?	NA	NA
What is the impact on equipment, including its use, function, serviceability, and so forth?	Less use of disinfectants on noncritical environmental surfaces and noncritical patient care equipment will prolong the useful life, serviceability and function of these surfaces and equipment due to the deleterious effects of the disinfectants. Score - 5	Prolonged use of germicides may have a deleterious effect on the equipment's use, function, and serviceability. Some germicides (alcohol and chlorine) can cause deterioration of rubber, plastics, and metals. Score - 1
What is the impact on internal physical systems?	NA	NA



Total Benefits Score = 23

Total Risk Score = 7



Meeting Actions: The team members scored each point as outlined, and concurred that there is no advantage to utilizing EPA-registered disinfectants at the recommended contact time of 10 minutes for disinfection of noncritical environmental surfaces and noncritical patient care equipment.

Regulatory Analysis: The Joint Commission

Joint Commission incorporates by reference the 2008 CDC Guideline for Disinfection and Sterilization in Healthcare Facilities, Recommendation 5K: Disinfect noncritical surfaces with an EPA-registered hospital disinfectant according to the label's safety precautions and use directions. Most EPA-registered hospital disinfectants have a label contact time of 10 minutes. However, many scientific studies have demonstrated the efficacy of hospital disinfectants against pathogens with a contact time of at least 1 minute. By law, the user must follow all applicable label instructions on EPA-registered products. If the user selects exposure conditions that differ from those on the EPA-registered product label, the user assumes liability for any injuries resulting from off-label use and is potentially subject to enforcement action under FIFRA. *Category II, IC.*

The Environmental Protection Agency

Under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA) requires all pesticides and disinfectants distributed and sold within the United States to be registered (licensed) by the EPA. Before the EPA may register a disinfectant/pesticide under FIFRA, they must demonstrate efficacy against the targeted organisms using the AOAC published methods and, according to specifications "will not generally cause unreasonable adverse effects on the environment". FIFRA also gives the EPA the ability regulate use through labeling, packaging, composition and disposal.

The Occupational Safety and Health Administration

1910.1030(d)(4)

Housekeeping --

1910.1030(d)(4)(i)

General. Employers shall ensure that the worksite is maintained in a clean and sanitary condition. The employer shall determine and implement an appropriate written schedule for cleaning and method of decontamination based upon the location within the facility, type of surface to be cleaned, type of soil present, and tasks or procedures being performed in the area.

1910.1030(d)(4)(ii)

All equipment and environmental and working surfaces shall be cleaned and decontaminated after contact with blood or other potentially infectious materials.

1910.1030(d)(4)(ii)(A)

Contaminated work surfaces shall be decontaminated with an appropriate disinfectant after completion of procedures; immediately or as soon as feasible when surfaces are overtly contaminated or after any spill of blood or other potentially infectious materials; and at the end of the work shift if the surface may have become contaminated since the last cleaning.

Management Data: No adverse or sentinel events or other healthcare-associated infection data suggests a problem with the current policy of using a 1 minute contact time for disinfection of non-critical environmental surfaces and noncritical patient care equipment.

Follow-up Plan: Incident reports will be monitored for any issues and if any identified, this policy will be reviewed.

Approval: The following leadership bodies have reviewed this risk assessment and adopted the position indicated. During this review, each leadership; body has concluded that this position is in the best interest of the patient.

Leadership Body	Date of Approval
Hospital Infection Control Committee	
Environmental Health and Safety Committee	

* References:

1. Rutala WA, White MS, Gergen MF, Weber DJ. 2006. Bacterial contamination of keyboards: Efficacy and functional impact of disinfectants. *Infect. Control Hosp. Epidemiol.* 27:372-7.
2. Rutala WA, Weber DJ. 2013. Disinfectants used for environmental disinfection and new room decontamination technology. *Am J Infect Control.* 41 (Supplement):S36-S41.
3. Rutala WA, Weber DJ. 2014. Selection of the ideal disinfectant. *Infect Control Hosp Epidemiol.* 35:855-865.
4. Rutala WA, Weber DJ. Monitoring and improving the effectiveness of surface cleaning/disinfection. *Am J Infect Control.* In press.
5. Huslage K, Rutala WA, Gergen MF, Sickbert-Bennett EE, Weber DJ. 2013. Microbial assessment of high, medium and low-touch hospital room surfaces. *Infect Control Hosp Epidemiol.* 34:211-212.
6. Sattar SA, Lloyd-Evans N, Springthorpe VS, Nair RC. Institutional outbreaks of rotavirus diarrhoea: potential role of fomites and environmental surfaces as vehicles for virus transmission. *J. Hyg. (Lond).* 1986;96:277-89.
7. Weber DJ, Rutala WA. Role of environmental contamination in the transmission of vancomycin-resistant enterococci. *Infect. Control Hosp. Epidemiol.* 1997;18:306-9.
8. Ward RL, Bernstein DI, Knowlton DR, et al. Prevention of surface-to-human transmission of rotaviruses by treatment with disinfectant spray. *J. Clin. Microbiol.* 1991;29:1991-6.
9. Tyler R, Ayliffe GA, Bradley C. Virucidal activity of disinfectants: studies with the poliovirus. *J. Hosp. Infect.* 1990;15:339-45.
10. Gwaltney JM, Jr., Hendley JO. Transmission of experimental rhinovirus infection by contaminated surfaces. *Am. J. Epidemiol.* 1982;116:828-33.



11. Sattar SA, Jacobsen H, Springthorpe VS, Cusack TM, Rubino JR. Chemical disinfection to interrupt transfer of rhinovirus type 14 from environmental surfaces to hands. *Appl. Environ. Microbiol.* 1993;59:1579-85.
12. Sattar SA, Jacobsen H, Rahman H, Cusack TM, Rubino JR. Interruption of rotavirus spread through chemical disinfection. *Infect. Control Hosp. Epidemiol.* 1994;15:751-6.
13. Rutala WA, Barbee SL, Aguiar NC, Sobsey MD, Weber DJ. Antimicrobial activity of home disinfectants and natural products against potential human pathogens. *Infect. Control Hosp. Epidemiol.* 2000;21:33-8.
14. Silverman J, Vazquez JA, Sobel JD, Zervos MJ. Comparative in vitro activity of antiseptics and disinfectants versus clinical isolates of *Candida* species. *Infect. Control Hosp. Epidemiol.* 1999;20:676-84.
15. Best M, Sattar SA, Springthorpe VS, Kennedy ME. Efficacies of selected disinfectants against *Mycobacterium tuberculosis*. *J. Clin. Microbiol.* 1990;28:2234-9.
16. Best M, Kennedy ME, Coates F. Efficacy of a variety of disinfectants against *Listeria* spp. *Appl. Environ. Microbiol.* 1990;56:377-80.
17. surfaces. *Appl. Environ. Microbiol.* 1990;56:3601-4.
18. Springthorpe VS, Grenier JL, Lloyd-Evans N, Sattar SA. Chemical disinfection of human rotaviruses: efficacy of commercially-available products in suspension tests. *J. Hyg. (Lond).* 1986;97:139-61.
19. Akamatsu T, Tabata K, Hironga M, Kawakami H, Uyeda M. Transmission of *Helicobacter pylori* infection via flexible fiberoptic endoscopy. *Am. J. Infect. Control* 1996;24:396-401.
20. Sattar SA, Springthorpe VS. Survival and disinfectant inactivation of the human immunodeficiency virus: a critical review. *Rev. Infect. Dis.* 1991;13:430-47.
21. Resnick L, Veren K, Salahuddin SZ, Tondreau S, Markham PD. Stability and inactivation of HTLV-III/LAV under clinical and laboratory environments. *JAMA* 1986;255:1887-91.
22. Weber DJ, Barbee SL, Sobsey MD, Rutala WA. The effect of blood on the antiviral activity of sodium hypochlorite, a phenolic, and a quaternary ammonium compound. *Infect. Control Hosp. Epidemiol.* 1999;20:821-7.
23. Rice EW, Clark RM, Johnson CH. Chlorine inactivation of *Escherichia coli* O157:H7. *Emerg. Infect. Dis.* 1999;5:461-3.
24. Rutala WA, Weber DJ, and the Healthcare Infection Control Practices Advisory Committee (HICPAC). Guideline for Disinfection and Sterilization in Healthcare Facilities, 2008. Pages 1-158. http://www.cdc.gov/hicpac/Disinfection_Sterilization/acknowledg.html. Accessed 15 March 2011.