
Best Practices in Disinfection of Noncritical Surfaces in the Healthcare Setting

William A. Rutala, Ph.D., M.P.H., C.I.C.

Director, Statewide Program for Infection Control and Epidemiology
and Professor of Medicine, University of North Carolina at Chapel
Hill, NC, USA

Former Director, Hospital Epidemiology, Occupational Health and
Safety, UNC Health Care, Chapel Hill, NC (1979-2017)

DISCLOSURES

2018

- Consultations
 - ASP (Advanced Sterilization Products), PDI
- Honoraria
 - PDI, ASP
- Scientific Advisory Board
 - Kinnos
- Grants
 - CDC, CMS

Disinfection of Noncritical Surfaces Bundle

NL Havill AJIC 2013;41:S26-30

- Develop policies and **procedures**
- Select cleaning and disinfecting **products**
- **Educate** staff-environmental services and nursing
- Monitor **compliance** (thoroughness of cleaning, product use) and feedback
- Implement **"no touch"** room decontamination technology and monitor compliance

www.disinfectionandsterilization.org

Environmental Contamination Leads to HAIs

Weber, Kanamori, Rutala. Curr Op Infect Dis 2016:29:424-431



- Evidence environment contributes
- Role-MRSA, VRE, *C. difficile*
- Surfaces are **contaminated**-~25%
- EIP **survive** days, weeks, months
- Contact with surfaces results in **hand contamination**
- Disinfection **reduces contamination**
- Disinfection (daily) **reduces HAIs**
- Rooms **not adequately cleaned**

Admission to Room Previously Occupied by Patient C/I with Epidemiologically Important Pathogen



- Results in the newly admitted patient having an increased risk of acquiring that pathogen by 39-353%
- For example, increased risk for *C. difficile* is 235% (11.0% vs 4.6%)
- Exposure to contaminated rooms confers a 5-6 fold increase in odds of infection, hospitals must adopt proven methods for reducing environmental contamination (Cohen et al. ICHE. 2018;39:541-546)

Objective

Institute Practices that Prevent All Infectious Disease
Transmission via Environment

Disinfection of Noncritical Surfaces Bundle

NL Havill AJIC 2013;41:S26-30

- Develop policies and procedures
- Select cleaning and disinfecting products
- Educate staff-environmental services and nursing
- Monitor compliance (thoroughness of cleaning, product use) and feedback
- Implement “no touch” room decontamination technology and monitor compliance

Disinfection of Noncritical Surfaces Bundle

- Develop policies and procedures
 - Environmental cleaning and disinfection is an integral part of preventing transmission of pathogens
 - In addition to identifying products and procedures, **ensure standardization of cleaning throughout the hospital**
 - ◆ Some units utilize ES to clean pieces of equipment (e.g., vital sign machines, IV pumps); some units use patient equipment, and some units utilize nursing staff.
 - ◆ Multidisciplinary group to create a standardized plan for cleaning patient rooms and pieces of patient equipment throughout the hospital

REVIEW THE “BEST” PRACTICES FOR CLEANING AND DISINFECTING

Cleaning and disinfecting is one-step with disinfectant-detergent. No pre-cleaning necessary unless spill or gross contamination. In many cases “best” practices not scientifically determined.

Blood Pressure Cuff

Non-Critical Patient Care Item



Surface Disinfection

Noncritical Patient Care

Rutala, Weber, HICPAC. CDC 2008. www.cdc.gov

- Disinfecting Noncritical Patient-Care Items
 - Process noncritical patient-care equipment with a **EPA-registered disinfectant** at the proper use dilution and a **contact time of at least 1 min**. *Category IB*
 - Ensure that the frequency for disinfecting noncritical patient-care surfaces be done minimally when **visibly soiled and on a regular basis** (such as after each patient use or once **daily** or once weekly). *Category IB*



Surface Disinfection

Environmental Surfaces

Rutala, Weber, HICPAC. CDC 2008. www.cdc.gov

- Disinfecting Environmental Surfaces in HCF
 - **Disinfect** (or clean) housekeeping surfaces (e.g., floors, tabletops) **on a regular basis** (e.g., **daily**, three times per week), **when spills occur, and when these surfaces are visibly soiled.**
Category IB
 - Use disinfectant for housekeeping purposes where: uncertainty exists as to the nature of the soil on the surfaces (blood vs dirt); or where uncertainty exists regarding the presence of multi-drug resistant organisms on such surfaces. *Category II*

Use of a Daily Disinfectant Cleaner Instead of a Daily Cleaner Reduced HAI Rates

Alfa et al. AJIC 2015.43:141-146

- Method: Improved hydrogen peroxide disposable wipe was used once per day for all high-touch surfaces to replace cleaner
- Result: When cleaning compliance was $\geq 80\%$, there was a significant reduction in cases/10,000 patient days for MRSA, VRE and *C. difficile*
- Conclusion: Daily use of disinfectant applied to environmental surfaces with a 80% compliance was superior to a cleaner because it resulted in significantly reduced rates of HAIs caused by *C. difficile*, MRSA, VRE

**It appears that not only is
disinfectant use important but
how often is important**

Daily disinfection vs clean when soiled

Daily Disinfection of High-Touch Surfaces

Kundrapu et al. ICHE 2012;33:1039

Daily disinfection of high-touch surfaces (vs cleaned when soiled) with sporicidal disinfectant (PA) in rooms of patients with CDI and MRSA reduced acquisition of pathogens on hands after contact with surfaces and of hands caring for the patient. **Daily disinfection less hand contamination.**




FIGURE 1. Effect of daily disinfection of high-touch environmental surfaces on acquisition of *Clostridium difficile* and methicillin-resistant *Staphylococcus aureus* (MRSA) on gloved hands of investigators after contact with the surfaces. A, Percentage of positive *C. difficile* cultures; B, mean number of *C. difficile* colony-forming units acquired; C, percentage of positive MRSA cultures; D, mean number of MRSA colony-forming units acquired.

EVIDENCE THAT ALL TOUCHABLE ROOM SURFACES ARE EQUALLY CONTAMINATED

TABLE 1. Precleaning and Postcleaning Bacterial Load Measurements for High-, Medium-, and Low-Touch Surfaces

Surface (no. of samples)	Mean CFUs/RODAC (95% CI)	
	Precleaning	Postcleaning
High (<i>n</i> = 40)	71.9 (46.5–97.3)	9.6 (3.8–15.4)
Medium (<i>n</i> = 42)	44.2 (28.1–60.2)	9.3 (1.2–17.5)
Low (<i>n</i> = 37)	56.7 (34.2–79.2)	5.7 (2.01–9.4)

NOTE. CFU, colony-forming unit; CI, confidence interval.

**Huslage K, Rutala W,
Gergen M, Sickbert-
Bennett S, Weber D
ICHE 2013;34:211-2**

Number of culture sites and prevalence of contamination with nosocomial pathogens in intensive care units (N=523)

Ward	Culture sites ^a			Prevalence of contamination
	HCWs' hands	Surfaces distant from patients	Surfaces close to patients	
A	3/10 (30%)	0/22 (0%)	6/25 (24.0%)	9/57 (15.8%)
B	2/9 (22.2%)	4/19 (21.1%)	5/48 (10.4%)	11/76 (14.5%)
C	2/10 (20%)	2/26 (7.7%)	7/49 (14.3%)	11/85 (12.9%)
D	1/9 (11.1%)	2/24 (8.2%)	7/45 (15.6%)	10/78 (12.8%)
E	0/5 (0%)	4/22 (18.2%)	3/30 (10%)	7/57 (12.3%)
F	1/10 (10%)	0/11 (0%)	4/31 (12.9%)	5/52 (9.6%)
G	0/3 (0%)	2/14 (14.3%)	0/20 (0%)	2/37 (5.4%)
H	1/10 (10%)	0/16 (0%)	1/55 (1.8%)	2/81 (2.5%)
Total	10/66 (15.2%)	14/154 (9.1%)	33/303 (10.9%)	57/523 (10.9%)

HCW, healthcare worker.

^a Number of contaminated samples/number of samples obtained.

**Willi I, Mayre A,
Kreidl P, et al.
JHI 2018;98:90-95**

ALL "TOUCHABLE" (HAND CONTACT) SURFACES SHOULD BE WIPED WITH DISINFECTANT

"High touch" objects only recently defined (no significant differences in microbial contamination of different surfaces) and "high risk" objects not epidemiologically defined. Cleaning and disinfecting is one-step with disinfectant-detergent. No pre-cleaning necessary unless spill or gross contamination.

Evaluation of Hospital Floors as a Potential Source of Pathogen Dissemination

Koganti et al. ICHE 2016. 37:1374; Deshpande et al. AJIC 2017. 45:336.

- Effective disinfection of contaminated surfaces is essential to prevent transmission of epidemiologically-important pathogens
- Efforts to improve disinfection focuses on touched surfaces
- Although floors contaminated, limited attention because not frequently touched
- Floors are a potential source of transmission because often contacted by objects that are then touched by hands (e.g., shoes, socks)
- Non-slip socks contaminated with MRSA, VRE (Mahida, J Hosp Infect. 2016;94:273)

Recovery of Nonpathogenic Viruses from Surfaces and Patients on Days 1, 2, and 3 After Inoculation of Floor Near Bed

Koganti et al. ICHE 2016. 37:1374

Variable	Day 1 (% Positive)	Day 2 (% Positive)	Day 3 (% Positive)
Patient Hands	40	63	43
Patient Footwear	100	100	86
High-touch surface <3ft	58	62	77
High-touch surface >3ft	40	68	34
Personal items	50	44	50
Adjacent room floor	NA	100	80
Adjacent room environment	NA	40	11
Nursing station	53	47	63
Portable equipment	33	23	100

Surfaces <3ft included bedrail, call button, telephone, tray table, etc; surfaces >3ft included side table, chair, IV pole, etc; personal-cell phones, books, clothing, wheelchairs; nurses station included computer keyboard, mouse, etc

Recovery of Nonpathogenic Viruses from Surfaces and Patients on Days 1, 2, and 3 After Inoculation of Floor Near Bed

Koganti et al. ICHE 2016. 37:1374

- Found that a nonpathogenic virus inoculated onto floors in hospital rooms disseminated rapidly to the footwear and hands of patients and to high-touch surfaces in the room
- The virus was also frequently found on high-touch surfaces in adjacent rooms and nursing stations
- Contamination in adjacent rooms in the nursing station suggest HCP contributed to dissemination after acquiring the virus during contact with surfaces or patients
- Studies needed to determine if floors are source of transmission

Disinfection of Noncritical Surfaces Bundle

- Develop policies and procedures
 - Standardize C/D patient rooms and pieces of equipment throughout the hospital
 - All touchable hand contact surfaces wiped with disinfection daily, when spills occur and when the surfaces are visibly soiled.
 - All noncritical medical devices should be disinfected daily and when soiled
 - Clean and disinfectant sink and toilet
 - Damp mop floor with disinfectant-detergent
 - If disinfectant prepared on-site, document correct concentration
 - Address treatment time/contact time for wipes and liquid disinfectants (e.g., treatment time for wipes is the kill time and includes a wet time via wiping as well as the undisturbed time).

Disinfection of Noncritical Surfaces Bundle

NL Havill AJIC 2013;41:S26-30

- Develop policies and procedures
- Select cleaning and disinfecting products
- Educate staff-environmental services and nursing
- Monitor compliance (thoroughness of cleaning, product use) and feedback
- Implement “no touch” room decontamination technology and monitor compliance

Effective Surface Decontamination

Product and Practice = Perfection

LOW-LEVEL DISINFECTION FOR NONCRITICAL EQUIPMENT AND SURFACES

Rutala, Weber. Infect Control Hosp Epidemiol. 2014;35:855-865


Exposure time \geq 1 min

Germicide	Use Concentration
Ethyl or isopropyl alcohol	70-90%
Chlorine	100ppm (1:500 dilution)
Phenolic	UD
Iodophor	UD
Quaternary ammonium (QUAT)	UD
QUAT with alcohol	RTU
Improved hydrogen peroxide (HP)	0.5%, 1.4%
Peracetic acid with HP (<i>C. difficile</i>)	UD

UD=Manufacturer's recommended use dilution; others in development/testing-electrolyzed water; polymeric guanidine; cold-air atmospheric pressure plasma (Boyce Antimicrob Res IC 2016. 5:10)

Microbiological Disinfectant Hierarchy

Rutala WA, Weber DJ, HICPAC. www.cdc.gov



MOST PREVALENT PATHOGENS CAUSING HAI

Rutala, Weber. Infect Control Hosp Epidemiol. 2014;35:855-865; Weiner et al ICHE 2016;37:1288

- Most prevalent pathogens causing HAI (easy to kill)
 - *E. coli* (15.4%)
 - *S. aureus* (11.8%)
 - *Klebsiella* (7.7%)
 - Coag neg Staph (7.7%)
 - *E. faecalis* (7.4%)
 - *P. aeruginosa* (7.3%)
 - *C. albicans* (6.7%)
 - *Enterobacter* sp. (4.2%)
 - *E. faecium* (3.7%)
- Common causes of outbreaks and ward closures (relatively hard to kill)
 - *C. difficile* spores
 - Norovirus
 - Rotavirus
 - Adenovirus

EFFECTIVENESS OF DISINFECTANTS AGAINST MRSA AND VRE

Rutala WA, et al. *Infect Control Hosp Epidemiol* 2000;21:33-38

TABLE 2
DISINFECTANT ACTIVITY AGAINST ANTIBIOTIC-SUSCEPTIBLE AND ANTIBIOTIC-RESISTANT BACTERIA

Product	Log ₁₀ Reductions							
	VSE		VRE		MSSA		MRSA	
	0.5 min	5 min	0.5 min	5 min	0.5 min	5 min	0.5 min	5 min
Vesphene IIse	>4.3	>4.3	>4.8	>4.8	>5.1	>5.1	>4.6	>4.6
Clorox	>5.4	>5.4	>4.9	>4.9	>5.0	>5.0	>4.6	>4.6
Lysol Disinfectant	>4.3	>4.3	>4.8	>4.8	>5.1	>5.1	>4.6	>4.6
Lysol Antibacterial	>5.5	>5.5	>5.5	>5.5	>5.1	>5.1	>4.6	>4.6
Vinegar	0.1	5.3	1.0	3.7	+1.1	+0.9	+0.6	2.3

Abbreviations: MRSA, methicillin-resistant *Staphylococcus aureus*; MSSA, methicillin-susceptible *S aureus*; VRE, vancomycin-resistant *Enterococcus*; VSE, vancomycin-susceptible *Enterococcus*. Data represent mean of two trials (n=2). Values preceded by ">" represent the limit of detection of the assay. Assays were conducted at a temperature of 20°C and a relative humidity of 45%. Results were calculated as the log of Nd/No, where Nd is the titer of bacteria surviving after exposure and No is the titer of the control.

Disinfection of Noncritical Surfaces Bundle

NL Havill AJIC 2013;41:S26-30

- Develop policies and procedures
- Select cleaning and disinfecting products
- Educate staff-environmental services and nursing
- Monitor compliance (thoroughness of cleaning, product use) and feedback
- Implement “no touch” room decontamination technology and monitor compliance

Disinfection of Noncritical Surfaces Bundle

- Develop policies and procedures
 - Standardize C/D patient rooms and pieces of equipment throughout the hospital
 - All touchable hand contact surfaces wiped with disinfection daily, when spills occur and when the surfaces are visibly soiled.
 - All noncritical medical devices should be disinfected daily and when soiled
 - Clean and disinfectant sink and toilet
 - Damp mop floor with disinfectant-detergent
 - If disinfectant prepared on-site, document correct concentration
 - Address treatment time/contact time for wipes and liquid disinfectants (e.g., treatment time for wipes is the kill time and includes a wet time via wiping as well as the undisturbed time).

Disinfection of Noncritical Surfaces Bundle

- Develop policies and procedures
 - Environmental cleaning and disinfection is an integral part of preventing transmission of pathogens
 - In addition to identifying products and procedures, **ensure standardization of cleaning throughout the hospital**
 - ◆ Some units utilize ES to clean pieces of equipment (e.g., vital sign machines, IV pumps); some units use patient equipment, and some units utilize nursing staff.
 - ◆ Multidisciplinary group to create a standardized plan for cleaning patient rooms and pieces of patient equipment throughout the hospital

Disinfection of Noncritical Surfaces Bundle

NL Havill AJIC 2013;41:S26-30


- Develop policies and procedures
- Select cleaning and disinfecting products
- Educate staff-environmental services and nursing
- Monitor compliance (thoroughness of cleaning, product use) and feedback
- Implement “no touch” room decontamination technology and monitor compliance

Effective Surface Decontamination

Product and Practice = Perfection

Thoroughness of Environmental Cleaning

Carling et al. ECCMID, Milan, Italy, May 2011



Practice* NOT Product

*surfaces not wiped

MONITORING THE EFFECTIVENESS OF CLEANING

Cooper et al. AJIC 2007;35:338

- Visual assessment-not a reliable indicator of surface cleanliness
- **ATP bioluminescence**-measures organic debris (each unit has own reading scale, <250-500 RLU)
- Microbiological methods-<2.5CFUs/cm²-pass; can be costly and pathogen specific
- **Fluorescent marker-transparent, easily cleaned, environmentally stable marking solution that fluoresces when exposed to an ultraviolet light** (applied by IP unbeknown to EVS, after EVS cleaning, markings are reassessed)

Thoroughness of Environmental Cleaning

Carling and Herwaldt. Infect Control Hosp Epidemiol 2017;38:960–965

Hospitals can improve their thoroughness of terminal room disinfection through fluorescent monitoring





FIGURE 4. A comparison of the results of the 3 previously published multisite studies compared with results from the Iowa project. White bars represent the average baseline TDCs and black bars represent the average final TDCs for sites that completed each study.

Percentage of Surfaces Clean by Different Measurement Methods


Rutala, Kanamori, Gergen, Sickbert-Bennett, Huslage, Weber. APIC Poster 2017.

Fluorescent marker is a useful tool in determining how thoroughly a surface is wiped and mimics the microbiological data better than ATP



Scatterplot of ATP Levels (less than 5000 RLU) and Standard Aerobic Counts (CFU/Rodac)

Rutala, Kanamori, Gergen, Sickbert-Bennett, Huslage, Weber. APIC 2017




There was no statistical correlation between ATP levels and standard aerobic plate counts.

Future Methods to Ensure Thoroughness

Future May Have Methods to Ensure Thoroughness Such as Colorized Disinfectant

Kang et al. J Hosp Infect 2017

Colorized disinfection – contact time compliance



0 min



2 min



4 min

- Color-fading time matched to disinfectant contact time --> enforces compliance
- Provides real-time feedback when disinfection is complete
- Trains staff on importance of contact time as they use the product

Colorized disinfection – improved coverage

Regular disinfectant wipes



Colorized wipes



- Increased visibility when disinfecting surfaces, fewer missed spots
- Real-time quality control that allows staff to monitor thoroughness of cleaning

Novel Chemical Additive That Colorizes Disinfectant to Improve Visualization of Surface Coverage

Mustapha et al . AJIC; 2018:48:191-121

By improving thoroughness will it reduce microbial contamination and reduce transmission?




Fig 1. (A) Percentage of sites correctly identified by personnel as having or not having bleach application when testing occurred within 30 seconds of application. (B) Image of a bed rail with application of bleach-plus-Highlight.

Disinfection of Noncritical Surfaces Bundle

NL Havill AJIC 2013;41:S26-30

- Develop policies and procedures
- Select cleaning and disinfecting products
- Educate staff-environmental services and nursing
- Monitor compliance (thoroughness of cleaning, product use) and feedback
- Implement “no touch” room decontamination technology and monitor compliance

These interventions (effective surface disinfection, thoroughness indicators) not enough to achieve consistent and high rates of cleaning/disinfection


No Touch

(supplements but do not replace surface cleaning/disinfection)

"NO TOUCH" APPROACHES TO ROOM DECONTAMINATION

(UV/VHP~20 microbicidal studies, 12 HAI reduction studies; will not discuss technology with limited data)

Weber, Kanamori, Rutala. Curr Op Infect Dis 2016;29:424-431; Weber, Rutala et al. AJIC; 2016:44: e77-e84; Anderson et al. Lancet 2017;389:805-14; Anderson et al. Lancet Infect Dis 2018;June 2018.



Enhanced Disinfection Leading to Reduction of Microbial Contamination and a Decrease in Patient Col/Infection

Anderson et al. Lancet 2017;289:805; Rutala et al. ICHE In press.

	Standard Method		Enhanced method	
	Quat	Quat/UV	Bleach	Bleach/UV
EIP (mean CFU per room) ^a	60.8	3.4	11.7	6.3
Reduction (%)		94	81	90
Colonization/Infection (rate) ^a	2.3	1.5	1.9	2.2
Reduction (%)		35	17	4

All enhanced disinfection technologies were significantly superior to Quat alone in reducing EIPs. Comparing the best strategy with the worst strategy (i.e., Quat vs Quat/UV) revealed that a reduction of 94% in EIP (60.8 vs 3.4) led to a 35% decrease in colonization/infection (2.3% vs 1.5%). Our data demonstrated that a decrease in room contamination was associated with a decrease in patient colonization/infection. First study which quantitatively described the entire pathway whereby improved disinfection decreases microbial contamination which in-turn reduced patient colonization/infection.

This technology ("no touch"-e.g., UV/HP) should be used (capital equipment budget) for terminal room disinfection (e.g., after discharge of patients on Contact Precautions).

Disinfection of Noncritical Surfaces Bundle

NL Havill AJIC 2013;41:S26-30

- Develop policies and procedures
- Select cleaning and disinfecting products
- Educate staff-environmental services and nursing
- Monitor compliance (thoroughness of cleaning, product use) and feedback
- Implement “no touch” room decontamination technology and monitor compliance

Our Responsibility to the Future

Institute Practices that Prevent All Infectious Disease
Transmission via Environment

Core Projects

- Inactivation of the human papillomavirus (HPV)
- Continuous room decontamination technologies for disinfection of the healthcare environment
 - Visible light disinfection through LEDs
 - Low concentration hydrogen peroxide
 - Persistent disinfectant that provides continuous disinfection action
 - ◆ Advantages and disadvantages of this technology
- Assess strategies to characterize/quantify environmental contamination to determine effectiveness of surface disinfection

Evaluation of A Persistent Surface Disinfectant Method

- Evaluation use the EPA "Protocol for Residual Self-Sanitizing Activity of Dried Chemical Residuals on Hard, Non-Porous Surfaces"
- Surfaces-glass, formica and SS
- Organisms- *S. aureus*, CRE and *C. auris*

Evaluation of A Persistent Surface Disinfectant Method



- Test method involves “wear” and re-inoculation of the test and control surfaces after
- Tester set to 5s for one pass
- Surface will undergo wear and re-inoculations over 24h
- Initial inoculation (10^5), apply disinfectant (dry overnight); 6 re-inoculations (10^3 , 30m dry), last inoculation (10^6)
- 24 passes (6 dry, 6 wet cycles)

EFFICACY OF A PERSISTENT CHEMICAL DISINFECTANT

Rutala WA, Gergen M, Sickbert-Bennett E, Anderson D, Weber D. Unpublished

- Methods: Surfaces were inoculated, treated with the novel disinfectant, allowed to dry, and then abraded using a standardized abrasion machine under multiple alternating wet and dry wipe conditions (N=12) interspersed with 6 re-inoculations. After 24 hours, the surface was re-inoculated a final time and ability of the disinfectant to kill >99.9% of 9 test microbes within 5min was measured on test surfaces (glass).
- **Persistent disinfectants may reduce or eliminate the problem of recontamination. Preliminary studies with a new persistent disinfectant are promising (4- 5 log₁₀ reduction in 5m over 24h).** When the novel disinfectant was compared to three other commonly used disinfectants using the same methodology with *S. aureus*, the mean log₁₀ reductions were: 4.4 (novel disinfectant); 0.9 (quat-alcohol); 0.2 (improved hydrogen peroxide); and 0.1 (chlorine).

Test Pathogen	Mean Log ₁₀ Reduction, 95% CI n=4
<i>S.aureus</i> *	4.4 (3.9, 5.0)
<i>S.aureus</i> (formica)	4.1 (3.8, 4.4)
<i>S.aureus</i> (stainless steel)	5.5 (5.2, 5.9)
VRE	≥4.5
<i>E.coli</i>	4.8 (4.6, 5.0)
<i>Enterobacter sp.</i>	4.1 (3.5, 4.6)
<i>Candida auris</i>	≥5.0
<i>K pneumoniae</i>	1.5 (1.4, 1.6)
CRE <i>E.coli</i>	3.0 (2.6, 3.4)
CRE <i>Enterobacter</i>	2.0 (1.6, 2.4)
CRE <i>K pneumoniae</i>	2.1 (1.8, 2.4)

Test surface glass unless otherwise specified

How Will We Prevent Infections Associated with the Environment?

- Implement evidence-based practices for surface disinfection
 - Evidence-based policies
 - Ensure use of safe and effective (against emerging pathogens such as *C. auris* and CRE) low-level disinfectants
 - Ensure thoroughness of cleaning (new thoroughness technology)
- Use “no touch” room decontamination technology proven to reduce microbial contamination on surfaces and reduction of HAIs at terminal/discharge cleaning
- Use new continuous room decontamination technology that continuously reduces microbial contamination

THANK YOU!

www.disinfectionandsterilization.org

